

AD\_\_\_\_\_

Award Number: DAMD17-02-1-0431

TITLE: Fatty Acid Synthase Inhibitor Cytotoxicity: Depletion of the Coenzyme-A Pool

PRINCIPAL INVESTIGATOR: Francis P. Kuhajda, M.D.

CONTRACTING ORGANIZATION: Johns Hopkins University School of Medicine  
Baltimore, MD 21205

REPORT DATE: April 2005

TYPE OF REPORT: Final, Revised

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE (DD-MM-YYYY) 01/04/05		2. REPORT TYPE Final, Revised		3. DATES COVERED (From - To) 1 Apr 2002 – 31 Mar 2005	
4. TITLE AND SUBTITLE  Fatty Acid Synthase Inhibitor Cytotoxicity: Depletion of the Coenzyme-A Pool				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER DAMD17-02-1-0431	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Francis P. Kuhajda, M.D.  E-Mail: fkuhajda@jhmi.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Johns Hopkins University School of Medicine Baltimore, MD 21205				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT: Inhibition of fatty acid synthase (FAS) with C75 in human cancer cells leads to cytotoxicity without evidence of DNA damage. Based on this and other biochemical observations, attention was focused on the cytoplasm as the site for the origin of C75 cytotoxicity to human cancer cells. We have shown that eukaryotic initiation factor 2 alpha (EIF2α), a key regulator of protein synthesis and the ER stress response, is involved in the cytotoxic mechanism of C75 against human breast cancer cells. Based on these observations, we synthesized a series of novel thiazolidinedione (TZD) compounds. TZD's likely induce cancer cell death through inhibition of translation initiation mediated by phosphorylation of eukaryotic initiation factor 2α (EIF2α) rendering it inactive. In summary, our novel TZD are cytotoxic to human colon cancer cells in vitro and xenografts in athymic mice. These compounds also phosphorylate EIF2α similar to C75. Based on these observations, novel class of potent TZD derivatives which may be useful for the treatment of a wide variety of human cancers.					
15. SUBJECT TERMS fatty acid synthase, thiazolidinediones, eukaryotic initiation factor 2α, HCT116 human colon cancer cells, endoplasmic reticulum stress response					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	16	19b. TELEPHONE NUMBER (include area code)

## Table of Contents

Introduction.....	4
Body.....	4
Key Research Accomplishments.....	7
Reportable Outcomes.....	7
Conclusions.....	7
References.....	7
Appendices.....	8

## **INTRODUCTION**

Initially, this proposal, written by Dr. Ellen Pizer at Johns Hopkins, sought to investigate coenzyme-A levels following inhibition of fatty acid synthase (FAS). Since we could not document consistent changes in coenzyme-A synthesis following fatty acid synthase inhibition, I suggested a change in the Statement of Work which was approved. FAS inhibition with C75 significantly inhibited protein synthesis in MCF7 cells within 4 hours. This reduction in protein synthesis is seen during apoptosis and is likely mediated by the ER stress response. Because of these findings, the role of eukaryotic initiation factor 2 alpha (EIF2 $\alpha$ ) was probed in regard to its effect of the regulation of protein translation. Interestingly, these studies have dovetailed with our overall interest in energy metabolism in cancer.

**BODY** Inhibition of fatty acid synthase (FAS) in human cancer cells leads to cytotoxicity without evidence of DNA damage. C75, the index FAS inhibitor that has been the focus of our mechanistic studies to determine how FAS inhibition triggers apoptosis, has been recently shown to also stimulate fatty acid oxidation (5, 6). It was conceivable that increased fatty acid oxidation could be the cause of cytotoxicity rather than FAS inhibition. In our recent paper, we have shown that FAS inhibition, not fatty acid oxidation stimulation is the trigger for apoptosis in human cancer cells (8). Given the dual effects of C75 on fatty acid synthesis and oxidation, we needed an additional compound with a different chemical scaffold to determine the relationship between the pharmacological manipulation of EIF2 $\alpha$  phosphorylation fatty acid synthesis inhibition, and cancer cell death before we could publish our results.

As part of our ongoing synthetic chemical effort, a series of novel thiazolidinedione (TZD) class of compounds were synthesized. During screening against human tumor cells, one particular compound, M133, inhibited fatty acid synthesis and was cytotoxic to MCF7 cells. The TZD class of compounds are well known ligands of peroxisome proliferators-activated receptor (PPAR)  $\gamma$  known to induce adipocyte differentiation and increase insulin sensitivity (4). This class of compounds have shown *in vitro* activity against lung (7), breast (1), colon (2), and prostate cancers (3), and against animal models of breast cancer (1). It was initially thought that TZD's were inducing apoptosis in human cancer cells by acting as PPAR $\gamma$  ligands. Recently, however, it was demonstrated that TZD's likely induce cancer cell death through inhibition of translation initiation mediated by phosphorylation of eukaryotic initiation factor 2 $\alpha$  (EIF2 $\alpha$ ) rendering it inactive (4).

We have now shown that similar to C75, M133 also rapidly inhibits fatty acid synthesis, phosphorylates EIF2, and has antitumor activity *in vitro* and *in vivo*. Using these data from disparate chemical scaffolds, we will now be able to demonstrate convincingly that inhibition of fatty acid synthesis leads to an ER stress response mediated through EIF2 $\alpha$  phosphorylation.


Recently, we have synthesized and tested a second novel TZD compound, M164 to determine if the anti-tumor effect of M133 was restricted to a single compound, potentially indicating an idiosyncratic effect, or common to this class of compounds. M164 also has significant anti-tumor activity both *in vitro* and *in vivo* demonstrating that these compound have potential as anti-tumor agents.

Through these series of experiments, we will outline the relationship FAS inhibition and the role of eukaryotic initiation factor 2 alpha (EIF2 $\alpha$ ) in the cytotoxic response using two novel TZD derivatives, M133 and M164.

***A. M133 inhibits fatty acid synthesis and is cytotoxic to human cancer cells.*** Using our well established method of [<sup>14</sup>C]acetate labeling to measure fatty acid synthesis, we have demonstrated that M133 inhibits fatty acid synthesis in MCF7 human breast cancer cells with an IC<sub>50</sub> of 15.4  $\mu$ g/ml within 2 h of drug application. As demonstrated by the *in vitro* XTT cytotoxicity assay, M133 has an IC<sub>50</sub> of 9.1 and 11.3  $\mu$ g/ml after 72h of drug


exposure against MCF7 and OVCAR3 human ovarian cancer cells. Both inhibition of fatty acid synthesis and cytotoxicity occur at concentrations  $\sim 30 \mu\text{M}$ .

***B. M133 induces EIF2 $\alpha$  phosphorylation.***



MCF7 cells were treated as indicated with M133 dissolved in DMSO. The DMSO concentration was kept below 0.1% and is included in the control wells. At concentrations well below the  $\text{IC}_{50}$ , M133 induces EIF2 $\alpha$  phosphorylation within 4h as shown by immunoblot (upper left panel) and persists at the 2  $\mu\text{g/ml}$  concentration at 24 h (upper right panel). The higher drug concentrations likely caused significant cell injury by 24 h. The lower panels show the expression of non-phosphorylated EIF2 $\alpha$  as a control.

***C. M133 causes significant growth inhibition of the HCT116 human colon cancer xenograft.***




In this pilot study, we utilized that fast growing, FAS expressing HCT116 human colon cancer cells grafted into the flanks of athymic mice. After 7 days of engraftment, with an average size of  $52 \text{ mm}^3$ , treatment with M133 began at 60 mg/kg, i.p., for the first two days, and 120 mg/kg on days 2,3, and 6. Control animals received 50  $\mu\text{l}$  DMSO vehicle. There were five animals in each group and tumors were measured with calipers in 3 dimensions. By day nine, there was a statistically significant difference between the controls and M133 treated animals. The study was terminated on day 10 due to compound limitations. Importantly, M133 does not cause the weight loss

seen with the C75 class of compounds because M133 does not induce fatty acid oxidation. Thus, we have achieved the separation of weight loss from cytotoxicity of fatty acid synthesis inhibitors.

**D. M121 has a significant anti-tumor effect both *in vitro* and *in vivo*.** M121 is a second novel TZD developed by our group. The rationale for developing and testing a second compound was to determine if the significant anti-tumor effects of M133 were idiosyncratic, or common to this class of compounds.

M164 was test against 3 human cancer cell lines *in vitro* with the following IC<sub>50</sub>'s: MCF7 (human breast cancer) 5.5 µg/ml, OVCAR-3 (human ovarian cancer) 3.7 µg/ml, and HCT116 (human colon cancer) 5.6 µg/ml. Fatty acid synthesis inhibition was measured by <sup>14</sup>Cactetate inhibition into total lipids in MCF7 cells with an IC<sub>50</sub> of ~ 30 µg/ml.

Using a similar treatment protocol to M133 described above, M164 has significant anti-tumor activity against the HCT116 xenograft in athymic mice as shown below.



Interestingly, compared to troglitazone against MCF7 xenografts, we achieved a superior result with 5 treatments over 9 days with both M133 and M164, than daily treatment at 1000 mg/kg for 9 weeks with troglitazone delivered orally (3). While this is an early study, we now have demonstrated that we have a TZD derivative that is likely significantly more potent *in vivo* than troglitazone. Since TZD's have been used in humans to treat diabetes, this compound may be similarly non-toxic, and if so, M133 or another derivative could be moved rapidly to the clinic.

**E. FAS inhibition activates AMP-activated protein kinase (AMPK).** We have recently demonstrated that FAS inhibition activates AMPK (9) (Zhou *et al*, 2007, Appendix I). Activated AMPK inhibits protein translation through inactivation of EIF as well. These data further indicate that inhibition of protein synthesis through a stress response or AMPK activation both demonstrate significant anti-tumor effects.

## **KEY RESEARCH ACCOMPLISHMENTS**

- A. Similar to C75, M133 inhibits fatty acid synthesis and is cytotoxic to human breast, colon, and ovarian cancer cells that express high levels of fatty acid synthesis.
- B. M133 does not cause significant weight loss similar to C75 because it does not stimulate fatty acid oxidation.
- C. M133 is potent against the HCT116 human colon cancer xenografts.
- D. M133 is a novel thiazolidinedione (TZD) derivative, a class of compounds used widely in the treatment of diabetes. As such, M133 could prove to be a relatively non-toxic compound with significant anti-tumor efficacy against a wide variety of human cancer.
- E. M164 is a second novel TZD derivative with significant *in vitro* and *in vivo* anti-tumor activity.

## **REPORTABLE OUTCOMES**

1. Publications: We are planning an abstract for the AACR meeting in 2008.
2. Degrees obtained: Minerva Hughes who synthesized these compound obtained her Ph. D. in Chemistry from the Johns Hopkins University.
3. Patents: We are in discussions over a patent for these compounds (composition of matter patent) for M133 and M164. The structures are patentable since they have not been previously synthesized.
4. Based on these data, we plan to apply for NIH funding during the following year in collaboration with the Townsend chemistry laboratory at JHU.

**CONCLUSIONS** Fatty acid synthase (FAS) is upregulated in most common human carcinomas. Inhibition of FAS has led to apoptosis in a variety of human cancer cell lines, and C75 has had significant anti-tumor activity in breast, prostate, colon, lung, and mesothelioma xenografts. Understanding the mechanism of cytotoxicity of FAS inhibitors is key to both the basic understanding of the biology and development of FAS as a drug target. Through this exploration of the mechanism of action of C75, we now have a novel class of potent TZD derivatives which may be useful for the treatment of a wide variety of human cancers.

## **REFERENCES**

1. **Elstner E, Muller C, Koshizuka K, Williamson EA, Park D, Asou H, Shintaku P, Said JW, Heber D, and Koeffler HP.** Ligands for peroxisome proliferator-activated receptor gamma and retinoic acid receptor inhibit growth and induce apoptosis of human breast cancer cells in vitro and in BNX mice. *Proc Natl Acad Sci U S A* 95: 8806-8811, 1998.
2. **Kitamura S, Miyazaki Y, Shinomura Y, Kondo S, Kanayama S, and Matsuzawa Y.** Peroxisome proliferator-activated receptor gamma induces growth arrest and differentiation markers of human colon cancer cells. *Jpn J Cancer Res* 90: 75-80, 1999.
3. **Kubota T, Koshizuka K, Williamson EA, Asou H, Said JW, Holden S, Miyoshi I, and Koeffler HP.** Ligand for peroxisome proliferator-activated receptor gamma (troglitazone) has potent antitumor effect against human prostate cancer both in vitro and in vivo. *Cancer Res* 58: 3344-3352, 1998.

4. **Palakurthi SS, Aktas H, Grubissich LM, Mortensen RM, and Halperin JA.** Anticancer effects of thiazolidinediones are independent of peroxisome proliferator-activated receptor gamma and mediated by inhibition of translation initiation. *Cancer Res* 61: 6213-6218, 2001.
5. **Thupari JN, Kim EK, Moran TH, Ronnett GV, and Kuhajda FP.** Chronic C75 Treatment of Diet-Induced Obese Mice Increases Fat Oxidation and Reduces Food Intake to Reduce Adipose Mass. *Am J Physiol Endocrinol Metab*, 2004.
6. **Thupari JN, Landree LE, Ronnett GV, and Kuhajda FP.** C75 increases peripheral energy utilization and fatty acid oxidation in diet-induced obesity. *Proc Natl Acad Sci U S A* 99: 9498-9502, 2002.
7. **Tsubouchi Y, Sano H, Kawahito Y, Mukai S, Yamada R, Kohno M, Inoue K, Hla T, and Kondo M.** Inhibition of human lung cancer cell growth by the peroxisome proliferator-activated receptor-gamma agonists through induction of apoptosis. *Biochem Biophys Res Commun* 270: 400-405, 2000.
8. **Zhou W, Simpson PJ, McFadden JM, Townsend CA, Medghalchi SM, Vadlamudi A, Pinn ML, Ronnett GV, and Kuhajda FP.** Fatty acid synthase inhibition triggers apoptosis during S phase in human cancer cells. *Cancer Res* 63: 7330-7337, 2003.
9. **Zhou W, Han WF, Landree LE, Thupari JN, Pinn ML, Bililign T, Kim EK, Vadlamudi A, Medghalchi SM, El Meskini R, Ronnett GV, Townsend CA, Kuhajda FP.** Fatty acid synthase inhibition activates AMP-activated protein kinase in SKOV3 human ovarian cancer cells. *Cancer Res* 67: 2964-2971, 2007.

**APPENDICES** . **Zhou W, Han WF, Landree LE, Thupari JN, Pinn ML, Bililign T, Kim EK, Vadlamudi A, Medghalchi SM, El Meskini R, Ronnett GV, Townsend CA, Kuhajda FP.** Fatty acid synthase inhibition activates AMP-activated protein kinase in SKOV3 human ovarian cancer cells. *Cancer Res* 67: 2964-2971, 2007.



# Fatty Acid Synthase Inhibition Activates AMP-Activated Protein Kinase in SKOV3 Human Ovarian Cancer Cells

Weibo Zhou,<sup>1</sup> Wan Fang Han,<sup>1</sup> Leslie E. Landree,<sup>2</sup> Jagan N. Thupari,<sup>1</sup> Michael L. Pinn,<sup>1</sup> Tsion Bililign,<sup>6</sup> Eun Kyoung Kim,<sup>2</sup> Aravinda Vadlamudi,<sup>7</sup> Susan M. Medghalchi,<sup>7</sup> Rajaa El Meskini,<sup>7</sup> Gabriele V. Ronnett,<sup>2,3</sup> Craig A. Townsend,<sup>6</sup> and Francis P. Kuhajda<sup>1,4,5</sup>

Departments of <sup>1</sup>Pathology, <sup>2</sup>Neuroscience, <sup>3</sup>Neurology, <sup>4</sup>Oncology, and <sup>5</sup>Biological Chemistry, The Johns Hopkins University School of Medicine; <sup>6</sup>Department of Chemistry, The Johns Hopkins University; and <sup>7</sup>FASgen, Inc., Baltimore, Maryland

## Abstract

**Fatty acid synthase (FAS), the enzyme responsible for the *de novo* synthesis of fatty acids, is highly expressed in ovarian cancers and most common human carcinomas. Inhibition of FAS and activation of AMP-activated protein kinase (AMPK) have been shown to be cytotoxic to human cancer cells *in vitro* and *in vivo*. In this report, we explore the cytotoxic mechanism of action of FAS inhibition and show that C93, a synthetic FAS inhibitor, increases the AMP/ATP ratio, activating AMPK in SKOV3 human ovarian cancer cells, which leads to cytotoxicity. As a physiologic consequence of AMPK activation, acetyl-CoA carboxylase (ACC), the rate-limiting enzyme of fatty acid synthesis, was phosphorylated and inhibited whereas glucose oxidation was increased. Despite these attempts to conserve energy, the AMP/ATP ratio increased with worsening cellular redox status. Pretreatment of SKOV3 cells with compound C, an AMPK inhibitor, substantially rescued the cells from C93 cytotoxicity, indicating its dependence on AMPK activation. 5-(Tetradecyloxy)-2-furoic acid, an ACC inhibitor, did not activate AMPK despite inhibiting fatty acid synthesis pathway activity and was not significantly cytotoxic to SKOV3 cells. This indicates that substrate accumulation from FAS inhibition triggering AMPK activation, not end-product depletion of fatty acids, is likely responsible for AMPK activation. C93 also exhibited significant antitumor activity and apoptosis against SKOV3 xenografts in athymic mice without significant weight loss or cytotoxicity to proliferating cellular compartments such as bone marrow, gastrointestinal tract, or skin. Thus, pharmacologic FAS inhibition selectively activates AMPK in ovarian cancer cells, inducing cytotoxicity while sparing most normal human tissues from the pleiotropic effects of AMPK activation. [Cancer Res 2007;67(7):2964–71]**

## Introduction

Although significant strides in the treatment and diagnosis of ovarian cancer have led to improved 5-year survival rates (1), ovarian cancer remains the leading cause of death from gynecologic

cancers (2). Fatty acid synthase (FAS), the enzyme responsible for the *de novo* synthesis of fatty acids, has emerged as a potential therapeutic target for human cancer. FAS catalyzes the condensation of malonyl-CoA and acetyl-CoA to produce long-chain fatty acids (3). High levels of FAS expression have been found in ovarian cancer (4, 5) and in most human solid tumors (6). The up-regulation of FAS expression in cancer cells has been linked to both mitogen-activated protein kinase and phosphatidylinositol 3-kinase pathways through the sterol regulatory binding element binding protein 1c (7–9). FAS protein expression denotes poor prognosis in breast and prostate cancer (10–15) and is found elevated in the blood of cancer patients (16, 17). Inhibition of FAS activity is selectively cytotoxic to human cancer cells *in vitro* and *in vivo* (13, 18–21) including human ovarian cancer xenografts (20, 22).

The mechanism linking inhibition of FAS activity to cancer cell death remains an active area of investigation. Because endogenously synthesized fatty acids in cancer cells are incorporated predominantly into phospholipids (23), it was hypothesized that FAS inhibition reduced available fatty acids for structural lipid synthesis in these cells. However, FAS inhibition induced apoptosis whereas pharmacologic inhibition of acetyl-CoA carboxylase (ACC), the rate-limiting enzyme of fatty acid synthesis, was far less toxic and actually protected cancer cells from FAS inhibition (18, 24). These findings implicated the accumulation of FAS substrates such as malonyl-CoA, rather than the depletion of end-product fatty acids, as contributing to cancer cell death. In addition to fatty acid synthesis, a number of other pathways have been found to modulate cytotoxicity following FAS inhibition including p53 (25, 26), HER2/*neu* (27, 28), and Akt (20, 29). Whereas these pathways may alter the response of cancer cells to FAS inhibition, the link between FAS enzyme inhibition and cancer cell death remains elusive.

C75, a synthetic mammalian FAS inhibitor (30), provided much of the *in vivo* evidence of the selective antitumor activity of pharmacologic FAS inhibition (13, 18, 19). Treatment, however, was limited by anorexia and weight loss (18, 31). While investigating the mechanism of action of C75-induced weight loss, we discovered that, in addition to FAS inhibition, C75 also stimulated carnitine palmitoyltransferase-1 (CPT-1) activity, leading to substantially increased fatty acid oxidation (32–35). CPT-1 is the rate-limiting transporter of long-chain acyl-CoAs into the mitochondria for oxidation. These studies raised the possibility that increased fatty acid oxidation could be responsible for at least part of the anticancer effect of C75. However, *in vitro* studies of fatty acid oxidation (24) and short interfering FAS RNA treatment of cancer cells (21) have established that FAS is the target of C75-induced cytotoxicity in cancer.

Although FAS seems to be the target of C75 for cancer cytotoxicity, the mechanism involved in the induction of anorexia

**Note:** Under a licensing agreement between FASgen and the Johns Hopkins University, F.P. Kuhajda and C.A. Townsend are entitled to a share of royalty received by the University on sales of products described in this article. F.P. Kuhajda and C.A. Townsend own and G.V. Ronnett has an interest in FASgen stock, which is subject to certain restrictions under University policy. The Johns Hopkins University, in accordance with its conflict of interest policies, is managing the terms of this arrangement.

**Requests for reprints:** Francis P. Kuhajda, Department of Pathology, Johns Hopkins Bayview Medical Center, 4940 Eastern Avenue, Baltimore, MD 21224. Phone: 410-550-0671; Fax: 410-550-0075; E-mail: fkuhajda@jhmi.edu.

©2007 American Association for Cancer Research.  
doi:10.1158/0008-5472.CAN-06-3439

involves the increase of fatty acid oxidation through CPT-1 stimulation, as well as the reduction of hypothalamic neuropeptide-Y expression (31). Further studies showed a pivotal role for AMP-activated protein kinase (AMPK) in the mechanism of C75-induced anorexia (36, 37). AMPK is activated in cells as a response to metabolic stresses that lead to the depletion of cellular ATP levels and thus the increase in AMP/ATP ratio (38, 39). The involvement of AMPK in the hypothalamic response to C75 (37) led to the hypothesis that the antitumor effect of FAS inhibition could also be mediated through changes in AMPK activation.

Thus, we and our collaborators sought to synthesize small-molecule FAS inhibitors that retain antitumor activity without inducing weight loss or stimulating fatty acid oxidation, which led to the development of C93 (40). In this study, we used human SKOV3 ovarian cancer cells to explore the role of AMPK in the cytotoxic mechanism of action of FAS inhibition using C93.

## Materials and Methods

**Cell lines, chemicals, and antibodies.** SKOV3 human ovarian cancer cells, obtained from the American Type Culture Collection (Manassas, VA), were cultured in McCoy's 5A medium with 10% FCS and 1% penicillin/streptomycin. C93 and compound C were obtained from FASgen, Inc. (Baltimore, MD). 5-(Tetradecyloxy)-2-furoic acid (TOFA) was synthesized by C.A. Townsend (Department of Chemistry, Johns Hopkins University, Baltimore, MD). C93, compound C, and TOFA were added to the cultures from 5 mg/mL stock solutions in 100% DMSO to give the indicated concentrations and the final DMSO concentration in cultures was  $\leq 0.2\%$ . Antibodies to pACC, pAMPK, AMPK, and  $\beta$ -actin were obtained from Cell Signaling (Danvers, MA). Mouse monoclonal antibodies to human FAS were obtained courtesy of FASgen.

To generate AMPK isoform-specific antibodies, rabbits were immunized with a peptide corresponding to amino acids 339 to 358 (DFY-LATSPDPSFLDDHHLTR) of rat AMPK $\alpha$ 1 or amino acids 352 to 366 (MDDSAMHIPPGLKPH) of  $\alpha$ 2 (41, 42). The keyhole limpet hemocyanin/GGG-conjugated peptides were injected to produce antibodies. The peptide-specific affinity-purified antibodies were used for SAMS peptide AMPK activity assay.

**Cell viability assay.** To measure the cytotoxicity of specific compounds against cancer cells,  $9 \times 10^3$  SKOV3 cells were plated per well onto 96-well plates. Following overnight culture, the compounds, dissolved in 100% DMSO, were added to the wells in 1- $\mu$ L volume at specified concentrations. Vehicle controls were run for each experiment. Each condition was run in triplicate.

After 24 or 72 h of incubation, cells were incubated for 4 h with the 2,3-bis[2-methoxy-4-nitro-5-sulfonylphenyl]-2H-tetrazolium-5-carboxanilide inner salt (XTT) reagent as per manufacturer's instructions (Cell Proliferation Kit II, Roche Diagnostics, Indianapolis, IN). Plates were read at  $A_{490}$  and  $A_{650}$  on a SpectraMax Plus Spectrophotometer (Molecular Devices, Sunnyvale, CA). Three wells containing the XTT reagent without cells served as the plate blank. XTT data were reported as  $A_{490} - A_{650}$ . Averages and SEs were computed using SOFTmax Pro software (Molecular Devices). The  $LC_{50}$  for each compound was defined as the concentration of drug leading to a 50% reduction in  $A_{490} - A_{650}$  compared with controls.  $LC_{50}$  was calculated by linear regression, plotting the FAS activity as percent of control versus drug concentrations. Linear regression, best-fit line,  $r^2$ , and 95% confidence intervals were determined using GraphPad Prism version 4.0 (GraphPad Software, San Diego, CA).

**Purification and assay of rat FAS.** Frozen livers from rats induced for FAS production were obtained from Pel-Freez Biologicals (Rogers, AR). Purification of FAS was adopted from Smith and Abraham (43) with minor modifications. An analogous protocol was used to isolate the human enzyme from a cultured ZR-75-1 breast cancer cell line provided by FASgen.

The specific activity and steady-state kinetic constants for overall FAS activity were similar to those reported in the literature for the rat (44) and

human (45) enzyme. Overall FAS activity was monitored spectrophotometrically by following the oxidation of NADPH at 240 nm [ $\epsilon = 6,220$  (mol/L) $^{-1}$  cm $^{-1}$ ] in the presence of the physiologic substrates malonyl-CoA and acetyl-CoA, as previously reported (43). Inhibition was monitored by varying concentrations of test compound in the overall FAS assay and  $IC_{50}$  values were determined by regression analysis (sigmoidal dose response) of the inhibition data (SigmaPlot, Point Richmond, CA).

**Fatty acid synthesis.** Fatty acid synthesis was measured by incorporation of [U- $^{14}$ C]acetate into lipids as described (18). Briefly, SKOV3 cells were cultured in 24-well plates at  $5 \times 10^4$  per well and incubated overnight. After the addition of drugs or vehicle alone as indicated, cells were pulse labeled with [U- $^{14}$ C]acetate, 1  $\mu$ Ci/well for 30 min to 2 h. Each condition was run in triplicate. Lipids were Folch extracted, counted for  $^{14}$ C, and data were expressed as percent of control.

**Fatty acid oxidation.** Fatty acid oxidation was measured by isolation of acid soluble products as described (46). Briefly,  $2.5 \times 10^5$  cells were plated onto 24-well plates. Following overnight culture, cells were incubated with drug for 60 min at 37°C for specified doses. The medium was removed and 250  $\mu$ L of reaction buffer [100  $\mu$ M/L [U- $^{14}$ C]palmitate (1.42  $\mu$ Ci), 200  $\mu$ M/L carnitine in serum-free medium with appropriate concentration of drug] were added for 30 min. The reaction was stopped with 50  $\mu$ L of 2.6 N HClO $_4$  and neutralized with 50  $\mu$ L of 4 N KOH. Cells were incubated at 60°C for 30 min and 75  $\mu$ L of 1 mol/L sodium acetate and 50  $\mu$ L of 3 N H $_2$ SO $_4$  were added. Cells were centrifuged at  $1,000 \times g$  for 7 min and extracted with 1:1 chloroform/methanol. The aqueous phase was counted for  $^{14}$ C. Each condition was run in triplicate; data were expressed as percent of control.

**Glucose oxidation.** Glucose oxidation was measured in SKOV3 cells as described (47). Briefly,  $10^5$  adherent SKOV3 cells were treated with C93 in Krebs-Ringer bicarbonate HEPES buffer containing 1% bovine serum albumin and 5 mmol/L glucose (47) at concentrations and times indicated. For the final 30 min of each treatment, 0.5  $\mu$ Ci/mL [U- $^{14}$ C]glucose was added. Reactions were stopped with 7% HClO $_4$  and 400  $\mu$ L of benzethonium hydroxide were injected in the center well. After 2 h at 37°C, complete oxidation was quantified by measuring the amount of  $^{14}$ C in the center well by liquid scintillation counting. Each condition was run in triplicate; data were expressed as percent of control.

**Western blot analysis.** Cells were homogenized in lysis buffer consisting of 50 mmol/L Tris-HCl (pH 7.5), 250 mmol/L sucrose, 5 mmol/L sodium pyrophosphate, 50 mmol/L NaF, 1 mmol/L EDTA, 1 mmol/L EGTA, 1 mmol/L DTT, 0.5 mmol/L benzamidin, 50  $\mu$ g/mL leupeptin, and 50  $\mu$ g/mL soybean trypsin inhibitor. SDS was added to a concentration of 0.2% and proteins were separated on 4% to 15% polyacrylamide gradient gels and transferred to a polyvinylidene difluoride membrane. Blots were probed with anti-pAMPK, anti-AMPK, anti-FAS, or anti- $\beta$ -actin. Samples for pACC immunoblots were run on separate 5% polyacrylamide gels.

**Measurement of adenine and pyridine nucleotides.** Both adenine and pyridine nucleotides (ATP, ADP, AMP, NAD $^+$ , NADH, NADP $^+$ , and NADPH) were measured concomitantly using high-performance liquid chromatography as described (48). Briefly,  $1 \times 10^6$  SKOV3 cells were plated onto 60-mm dishes and treated as described. Pyridine and adenine nucleotides were extracted with alkali before high-performance liquid chromatography analysis. Cell extracts and purified nucleotide controls (Sigma) were separated and quantified using an Agilent 1100 running ChemStation software as follows: 50  $\mu$ L of the supernatants were loaded onto an LC-18 column and absorbance at 254 nm was measured. The two eluants were buffer A: 0.1 mol/L KH $_2$ PO $_4$ , 8 mmol/L tetrabutylammonium hydrogen sulfate, pH 6.0; and buffer B: 70% buffer A in methanol. The following gradients were carried out: 0 to 2.5 min, 100% A; 2.5 to 5.0 min, 80% A; 5.0 to 10.0 min, 60% A; 10.0 to 13.0 min, 0% A (100% B), for 5 min. The flow rate was 1.5 mL/min. Data were expressed as ratios of redox couples or as absorbance at 254 nm.

**Measurement of AMPK activity.** AMPK activity was measured by carrying out the SAMS peptide assay as described (36, 49). SKOV3 cells were plated onto six-well dishes and three wells were used per condition. AMPK $\alpha$ 1 or AMPK $\alpha$ 2 isoforms were immunoprecipitated in the presence of isoform-specific antibodies coupled to protein A/G beads (Santa Cruz



Biotechnology, Santa Cruz, CA). Immunoprecipitates were washed and kinase activity was assessed by incorporation of  $^{32}\text{P}$  into the synthetic SAMS peptide substrate (Princeton Biomolecules, Langhorne, PA). Each sample was corrected for protein concentration and triplicate assays were reported as percent of control.

**Animal studies.** All animal experiments were done in accordance with guidelines on animal care and use established by the Johns Hopkins University School of Medicine Institutional Animal Care and Use Committee. Female athymic nude BALB/c mice (8–10 weeks) were purchased from Harlan Laboratories (Indianapolis, IN) and housed in a pathogen-free facility. Animals were treated once or twice daily with i.p. injections of C93, with 50 mg/kg dissolved in 50  $\mu\text{L}$  of 100% DMSO, or with vehicle alone as specified. For i.p. xenografts,  $5 \times 10^7$  SKOV3 cells were injected. After seven days, C93 was administered i.p. at 50 mg/kg daily in DMSO every fifth day for 4 weeks. Wet tumor weight was obtained at necropsy. For flank xenografts,  $2 \times 10^6$  SKOV3 cells were injected and treatment commenced when tumors reached 100  $\text{mm}^3$ . C93 was administered i.p. at 50 mg/kg twice a day in 100% DMSO on each day except days 6 and 13. Tumor volume and animal weights were measured daily.

## Results

**C93 inhibits FAS and is cytotoxic to SKOV3 cells but does not increase fatty acid oxidation or induce weight loss.** C93 emerged from a screen of potential inhibitors assayed against homogenous FAS isolated from rat liver and ZR-75-1 human breast cancer cells. The development of C93 accomplished the pharmacologic elimination of CPT-1 stimulation while preserving FAS inhibition. Thus, C93 does not substantially affect weight loss or feeding behavior as seen with C75 (40, 50). Highlighted in Table 1 are key features of C93 compared with C75 and TOFA, an inhibitor of ACC (51). Both C93 and C75 inhibit FAS, but C93 is 2-fold more effective in pathway inhibition and >3-fold more toxic to SKOV3 cells. Importantly, C93 does not stimulate CPT-1 or increase fatty acid oxidation as does C75. As a result, C93 induced a 1.4% weight loss in lean BALB/c mice with a single i.p. 60 mg/kg dose, compared with a 15% weight loss with a single i.p. 30 mg/kg dose of C75. Similar to our previous report (18), ACC inhibition with TOFA was less cytotoxic than FAS inhibition. TOFA, although >25-fold more effective in pathway inhibition than C93 in SKOV3 cells, was 16-fold less effective as a cytotoxic agent against SKOV3 cells compared with C93. Thus, inhibition of the fatty acid synthesis pathway by FAS is substantially more efficacious compared with inhibition by the physiologic rate-limiting enzyme ACC.

**C93 rapidly inhibits fatty acid synthesis and activates AMPK.** To establish the time course of events following C93 treatment of SKOV3 cells, we first determined how rapidly C93 inhibited fatty acid synthesis by measuring the incorporation of [ $^{14}\text{C}$ ]acetate into

lipids (Fig. 1A). Initially, C93 inhibited fatty acid synthesis pathway activity by ~60% relative to vehicle control by 30 min posttreatment. By 160 min posttreatment, fatty acid synthesis activity decreased to 40% of the control. We next measured ATP and AMP levels to assess cellular energy status over the same time course. Using the same concentrations of C93 and culture conditions, a significant elevation in the AMP/ATP ratio occurred within 30 min of C93 treatment (Fig. 1B). The AMP/ATP ratio continued to increase to >2-fold compared with control at 180 min posttreatment. Concomitant with the increase in the AMP/ATP ratio, C93 treatment enhanced AMPK phosphorylation detected by immunoblot within 30 min of treatment persisting through 180 min (Fig. 1C). AMPK phosphorylation was accompanied by phosphorylation of one of its target proteins, ACC, detected 120 min posttreatment. The activation of AMPK has been shown to phosphorylate and inhibit ACC, thus reducing fatty acid synthesis pathway activity, thereby preserving ATP (52). Thus, ACC phosphorylation may be responsible for the continued reduction in FAS pathway activity at 160 min post C93 treatment seen in Fig. 1A. Recently, AMPK activation has been reported to reduce FAS expression (53). Whereas changes in FAS expression would be unlikely to account for the rapid reduction in pathway activity, nonetheless, an immunoblot showed no change in FAS protein expression compared with the  $\beta$ -actin control during the course of C93 treatment (Fig. 1C).

Increased AMPK phosphorylation denotes increased AMPK activity. To confirm the biological significance of the AMPK phosphorylation and directly measure AMPK activity, SAMS peptide activity assays were done (Fig. 1D). In the inset, immunoblots of SKOV3 cells showed predominance of the  $\alpha 2$  isoform over the  $\alpha 1$  isoform. Using antibodies selective for the  $\alpha 2$  isoform, an ~150% increase in AMPK activity was noted at 30 min following C93 treatment, which continued to increase to >200% of control by 120 min posttreatment. Taken together, these data strongly suggest that C93 treatment rapidly activates AMPK in SKOV3 cells.


**C93 increases glucose oxidation while reducing both  $\text{NAD}^+$ /NADH and  $\text{NADP}^+$ /NADPH ratios.** Whereas AMPK activation reduces the activity of anabolic pathways to preserve energy expenditure, it concomitantly increases the activity of catabolic pathways in an attempt to replenish ATP levels. Figure 2A shows that following C93 treatment, SKOV3 cells significantly increased glucose oxidation compared with controls within 60 min of treatment. However, despite the increased glucose oxidation noted at 60 min posttreatment, the AMP/ATP ratio did not improve (Fig. 1B). In addition to the increasing AMP/ATP ratio, Fig. 2B

**Table 1.** Comparison of C93, C75, and TOFA

Drug	Target	FAS inhibition $\text{IC}_{50}$ ( $\mu\text{g}/\text{mL}$ )	Pathway inhibition $\text{IC}_{50}$ ( $\mu\text{g}/\text{mL}$ )	SKOV3 XTT $\text{LC}_{50}$ ( $\mu\text{g}/\text{mL}$ )	CPT-1 stimulation	FAO stimulation $\text{SC}_{150}$ ( $\mu\text{g}/\text{mL}$ )	Maximum weight loss [% (i.p. dose)]
C93	FAS	30	$5.5 \pm 0.5$	$2.6 \pm 0.3$	None	None	1.4 (60 mg/kg)
C75	FAS/CPT-1	10	$10.8 \pm 2.7$	$11.0 \pm 2.3$	300% at 20 $\mu\text{g}/\text{mL}$	$1.7 \pm 2.0$	15.0 (30 mg/kg)
TOFA	ACC	NA	$0.20 \pm 0.024$	$42.4 \pm 1.8$	None	None	0 (60 mg/kg)

Abbreviations: FAO, fatty acid oxidation;  $\text{SC}_{150}$ , stimulatory concentration 150; NA, not applicable.

**Figure 1.** C93 inhibits fatty acid synthesis and increases the AMP/ATP ratio, activating AMPK in SKOV3 cells. **A**, C93 inhibited fatty acid synthesis within 30 min of treatment with further inhibition occurring 160 min posttreatment (two-tailed *t* tests). **B**, the AMP/ATP ratio increased within 30 min following C93 treatment ( $P < 0.0001$ ). The overall *P* value was calculated by one-way ANOVA analysis; *P* values for all individual time points were calculated with time 0 as the control value using Dunnett's posttest. **C**, immunoblot showing AMPK and ACC phosphorylation 30 and 120 min following C93, respectively. There was no change in FAS expression. **D**, following C93 treatment, AMPK activity, as measured by SAMS peptide assay, increased as compared with vehicle control (two-tailed *t* tests). AMPK activity paralleled AMPK phosphorylation as shown in (C). Immunoblots showed that SKOV3 cells predominantly express the  $\alpha 2$  AMPK isoform (insets; \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ ; GraphPad Software).



shows that following C93 treatment, the NAD<sup>+</sup>/NADH ratio declined, indicating a failure of the cell to maintain a positive redox balance.

FAS uses the reductive power of NADPH in the synthesis of fatty acids with the oxidation of 14 mol of NADPH per mole of fatty acid synthesized. Thus, inhibition of FAS might be expected to increase, at least transiently, NADPH levels. Figure 2C shows that NADPH levels indeed increased within 5 min following C93 treatment without a substantial change in the NADPH/NADP<sup>+</sup> ratio (Fig. 2D). After 60 min, however, the NADPH/NADP<sup>+</sup> ratio begins to decline, which may reflect the overall waning in reductive biosynthesis activity that follows AMPK activation (52).

**Compound C rescues C93-induced cytotoxicity.** If AMPK activation contributes to C93 cytotoxicity, blocking AMPK activation with compound C, an AMPK inhibitor (37, 54), should blunt the cytotoxic effect of C93 treatment. Figure 3A represents three separate experiments in which SKOV3 cells were treated with C93 alone, C93 combined with a 1-hour pretreatment with compound C, or compound C alone. Compound C pretreatment substantially reduced the cytotoxicity of C93 assayed 24 h post-treatment. Treatment with compound C alone at the same concentration had no significant cytotoxic effect on SKOV3 cells. These data suggest that AMPK activation plays a substantial role in the cytotoxic mechanism of action of C93.

**TOFA does not activate AMPK and is not toxic to SKOV3 cells at a concentration that substantially inhibits fatty acid synthesis.** Because ACC inhibition has been shown to be less cytotoxic to human cancer cells than inhibition of fatty acid synthesis at FAS (18), we hypothesized that TOFA treatment would not increase AMPK phosphorylation. In Fig. 3B, TOFA treatment at 5  $\mu$ g/mL, a concentration that dramatically reduced fatty acid synthesis (Table 1), did not substantially affect AMPK phosphor-

ylation. In contrast, 5-aminoimidazole-4-carboxamide-1- $\beta$ -D-ribofuranoside (AICAR), an activator of AMPK, caused substantial AMPK phosphorylation in SKOV3 cells at 1 mmol/L (Fig. 3B, lane A).

Because TOFA does not activate AMPK, we next compared TOFA with C93 and AICAR in a cytotoxicity assay (Fig. 3C). After 24 h, C93 reduced the viability of SKOV3 cells to 75% of vehicle control, whereas TOFA, at a concentration that substantially inhibits fatty acid synthesis pathway activity, did not significantly affect cell viability. AICAR, an activator of AMPK, was also substantially cytotoxic. Thus, C93 and AICAR, which both activate AMPK, were cytotoxic to SKOV3 cells, whereas TOFA was not. These data further substantiate that (a) AMPK activation is cytotoxic to human cancer cells (55); (b) inhibition of ACC with TOFA does not lead to increased AMPK phosphorylation, consistent with previous experiments using cultured neurons (36); and (c) inhibition of fatty acid synthesis does not activate AMPK.

**C93 inhibits the growth of the SKOV3 xenograft.** Whereas C93 is effective as a cytotoxic agent against SKOV3 cells *in vitro*, we did a pilot studies of C93 to assess the antitumor activity of C93 in SKOV3 xenograft-bearing mice. In the SKOV3 i.p. xenograft model (ref. 20; Fig. 4A), the five informative C93-treated animals had an average tumor volume of ~14 g compared with 110 g for the vehicle controls. Three of the treated animals had no identifiable residual tumor. No significant weight loss was identified. There were peritoneal adhesions in the C93-treated animals bearing i.p. xenografts. In this group, three animals were lost due to gastrointestinal trauma during i.p. injections.

Figure 4B illustrates the antitumor effect of C93 on a s.c. xenograft of SKOV3 cells in athymic mice. C93 inhibited tumor growth 3-fold compared with vehicle controls ( $P < 0.0001$ , repeated measures two-way ANOVA analysis; GraphPad Prism version 4.00 for Windows). At

- Tondo U. Expression of fatty acid synthase (FAS) as a predictor of recurrence in stage I breast carcinoma patients. *Cancer* 1996;77:474–82.
15. Milgram LZ, Witters LA, Pasternack GR, Kuhajda FP. Enzymes of the fatty acid synthesis pathway are highly expressed in *in situ* breast carcinoma. *Clin Cancer Res* 1997;3:2115–20.
  16. Wang Y, Kuhajda FP, Sokoll LJ, Chan DW. Two-site ELISA for the quantitative determination of fatty acid synthase. *Clin Chim Acta* 2001;304:107–15.
  17. Wang YY, Kuhajda FP, Cheng P, et al. A new model ELISA, based on two monoclonal antibodies, for quantification of fatty acid synthase. *J Immunoassay Immunochem* 2002;23:279–92.
  18. Pizer E, Thupari J, Han W, et al. Malonyl-Coenzyme-A is a potential mediator of cytotoxicity induced by fatty acid synthase inhibition in human breast cancer cells and xenografts. *Cancer Res* 2000;60:213–8.
  19. Gabrielson EW, Pinn ML, Testa JR, Kuhajda FP. Increased fatty acid synthase is a therapeutic target in mesothelioma. *Clin Cancer Res* 2001;7:153–7.
  20. Wang HQ, Altomare DA, Skele KL, et al. Positive feedback regulation between AKT activation and fatty acid synthase expression in ovarian carcinoma cells. *Oncogene* 2005;24:3574–82.
  21. De Schrijver E, Brusselmans K, Heyns W, Verhoeven G, Swinnen JV. RNA interference-mediated silencing of the fatty acid synthase gene attenuates growth and induces morphological changes and apoptosis of LNCaP prostate cancer cells. *Cancer Res* 2003;63:3799–804.
  22. Pizer E, Wood F, Heine H, Romantsev F, Pasternack G, Kuhajda F. Inhibition of fatty acid synthesis delays disease progression in a xenograft model of ovarian cancer. *Cancer Res* 1996;56:1189–93.
  23. Kuhajda FP, Jenner K, Wood FD, et al. Fatty acid synthase: a potential selective target for antineoplastic therapy. *Proc Natl Acad Sci U S A* 1994;91:6379–83.
  24. Zhou W, Simpson PJ, McFadden JM, et al. Fatty acid synthase inhibition triggers apoptosis during S-phase in human cancer cells. *Cancer Res* 2003;63:7330–7.
  25. Li JN, Gorospe M, Chrest FJ, et al. Pharmacological inhibition of fatty acid synthase activity produces both cytostatic and cytotoxic effects modulated by p53. *Cancer Res* 2001;61:1493–9.
  26. Menendez JA, Lupu R. RNA interference-mediated silencing of the p53 tumor-suppressor protein drastically increases apoptosis after inhibition of endogenous fatty acid metabolism in breast cancer cells. *Int J Mol Med* 2005;15:33–40.
  27. Menendez JA, Vellon L, Mehmi I, et al. Inhibition of fatty acid synthase (FAS) suppresses HER2/neu (erbB-2) oncogene overexpression in cancer cells. *Proc Natl Acad Sci U S A* 2004;101:10715–20.
  28. Kumar-Sinha C, Ignatoski KW, Lippman ME, Ethier SP, Chinnaiyan AM. Transcriptome analysis of HER2 reveals a molecular connection to fatty acid synthesis. *Cancer Res* 2003;63:132–9.
  29. Alli PM, Pinn ML, Jaffee EM, McFadden JM, Kuhajda FP. Fatty acid synthase inhibitors are chemopreventive for mammary cancer in neu-N transgenic mice. *Oncogene* 2005;24:39–46.
  30. Kuhajda FP, Pizer EP, Li JL, Mani NS, Frehywot GL, Townsend CA. Synthesis and anti-tumor activity of a novel inhibitor of fatty acid synthase. *Proc Natl Acad Sci U S A* 2000;97:3450–4.
  31. Lofthus TM, Jaworsky DE, Frehywot GL, et al. Reduced food intake and body weight in mice treated with fatty acid synthase inhibitors. *Science* 2000;288:2379–81.
  32. Thupari JN, Kim EK, Moran TH, Ronnett GV, Kuhajda FP. Chronic C75 treatment of diet-induced obese mice increases fat oxidation and reduces food intake to reduce adipose mass. *Am J Physiol Endocrinol Metab* 2004;287:E97–104.
  33. Thupari JN, Landree LE, Ronnett GV, Kuhajda FP. C75 increases peripheral energy utilization and fatty acid oxidation in diet-induced obesity. *Proc Natl Acad Sci U S A* 2002;99:9488–502.
  34. Nicot C, Napal L, Relat J, et al. C75 activates malonyl-CoA sensitive and insensitive components of the CPT system. *Biochem Biophys Res Commun* 2004;325:660–4.
  35. Yang N, Kays JS, Skillman TR, Burris L, Seng TW, Hammond C. C75 activates carnitine palmitoyltransferase-1 in isolated mitochondria and intact cells without displacement of bound malonyl CoA. *J Pharmacol Exp Ther* 2005;312:127–33.
  36. Landree LE, Hanlon AL, Strong DW, et al. C75, a fatty acid synthase inhibitor, modulates AMP-activated protein kinase to alter neuronal energy metabolism. *J Biol Chem* 2004;279:3817–27.
  37. Kim EK, Miller I, Aja S, et al. C75, a fatty acid synthase inhibitor, reduces food intake via hypothalamic AMP-activated protein kinase. *J Biol Chem* 2004;279:19970–6.
  38. Hardie DG, Carling D, Carlson M. The AMP-activated/SNF1 protein kinase subfamily: metabolic sensors of the eukaryotic cell? *Annu Rev Biochem* 1998;67:821–55.
  39. Kemp BE, Mitchellhill KI, Stapleton D, Michell BJ, Chen ZP, Witters LA. Dealing with energy demand: the AMP-activated protein kinase. *Trends Biochem Sci* 1999;24:22–5.
  40. McFadden JM, Medghalchi SM, Thupari JN, et al. Application of a flexible synthesis of (5R)-thiolactomycin to develop new inhibitors of type I fatty acid synthase. *J Med Chem* 2005;48:946–61.
  41. Woods A, Cheung PC, Smith FC, et al. Characterization of AMP-activated protein kinase  $\beta$  and  $\gamma$  subunits. Assembly of the heterotrimeric complex *in vitro*. *J Biol Chem* 1996;271:10282–90.
  42. Xing Y, Musi N, Fujii N, et al. Glucose metabolism and energy homeostasis in mouse hearts overexpressing dominant negative  $\alpha 2$  subunit of AMP-activated protein kinase. *J Biol Chem* 2003;278:28372–7.
  43. Smith S, Abraham S. Fatty acid synthase from lactating rat mammary gland. *Methods Enzymol* 1975;35:65–74.
  44. Aprahamian SA, Arslanian MJ, Wakil SJ. Comparative studies on the kinetic parameters and product analyses of chicken and rat liver and yeast fatty acid synthetase. *Comp Biochem Physiol B Biochem Mol Biol* 1982;71:577–82.
  45. Jayakumar A, Tai MH, Huang WY, et al. Human fatty acid synthase: properties and molecular cloning. *Proc Natl Acad Sci U S A* 1995;92:8695–9.
  46. Watkins PA, Ferrell EV, Jr, Pedersen JL, Hoefler G. Peroxisomal fatty acid  $\beta$ -oxidation in HepG2 cells. *Arch Biochem Biophys* 1991;289:329–36.
  47. Rubi B, Antinozzi PA, Herrero L, et al. Adenovirus-mediated overexpression of liver carnitine palmitoyltransferase I in INSIE cells: effects on cell metabolism and insulin secretion. *Biochem J* 2002;364:219–26.
  48. Stocchi V, Cucchiari L, Magnani M, Chiarantini L, Palma P, Crescentini G. Simultaneous extraction and reverse-phase high-performance liquid chromatographic determination of adenine and pyridine nucleotides in human red blood cells. *Anal Biochem* 1985;146:118–24.
  49. Witters LA, Kemp BE. Insulin activation of acetyl-CoA carboxylase accompanied by inhibition of the 5'-AMP-activated protein kinase. *J Biol Chem* 1992;267:2864–7.
  50. McFadden JM, Frehywot GL, Townsend CA. A flexible route to (5R)-thiolactomycin, a naturally occurring inhibitor of fatty acid synthesis. *Org Lett* 2002;4:3859–62.
  51. Halvorson DL, McCune SA. Inhibition of fatty acid synthesis in isolated adipocytes by 5-(tetradecyloxy)-2-furoic acid. *Lipids* 1984;19:851–6.
  52. Hardie DG, Hawley SA. AMP-activated protein kinase: the energy charge hypothesis revisited. *BioEssays* 2001;23:1112–9.
  53. Woods A, Azzout-Marniche D, Foretz M, et al. Characterization of the role of AMP-activated protein kinase in the regulation of glucose-activated gene expression using constitutively active and dominant negative forms of the kinase. *Mol Cell Biol* 2000;20:6704–11.
  54. Al-Khalili L, Chibalin AV, Yu M, et al. MEK2 activation in differentiated primary human skeletal muscle cultures requires coordinated involvement of parallel pathways. *Am J Physiol Cell Physiol* 2004;286:C1410–6.
  55. Rattan R, Giri S, Singh AK, Singh I. 5-Aminoimidazole-4-carboxamide-1- $\beta$ -D-ribofuranoside inhibits cancer cell proliferation *in vitro* and *in vivo* via AMP-activated protein kinase (AMPK). *J Biol Chem* 2005;280:39582–93.
  56. Pizer E, Pflug B, Bova G, Han W, Udan M, Nelson J. Increased fatty acid synthase as a therapeutic target in androgen-independent prostate cancer progression. *Prostate* 2001;47:102–10.
  57. Aja S, McFadden JM, Aplasca A, et al. Intracerebroventricular administration of either C89b, a stimulator of carnitinepalmitoyl-transferase-1 (CPT-1s), or cerulenin, an inhibitor of fatty acid synthase (FAS), reduces food intake and body weight in mice. *Appetite* 2006;46:339.
  58. Hardie DG, Carling D. The AMP-activated protein kinase-fuel gauge of the mammalian cell? *Eur J Biochem* 1997;246:259–73.
  59. Abbud W, Habinowski S, Zhang JZ, et al. Stimulation of AMP-activated protein kinase (AMPK) is associated with enhancement of GLUT1-mediated glucose transport. *Arch Biochem Biophys* 2000;380:347–52.
  60. Kurth-Kraczek EJ, Hirshman MF, Goodyear LJ, Winder WW. 5' AMP-activated protein kinase activation causes GLUT4 translocation in skeletal muscle. *Diabetes* 1999;48:1667–71.
  61. Marsin AS, Bertrand L, Rider MH, et al. Phosphorylation and activation of heart PFK-2 by AMPK has a role in the stimulation of glycolysis during ischaemia. *Curr Biol* 2000;10:1247–55.
  62. Winder WW, Hardie DG. Inactivation of acetyl-CoA carboxylase and activation of AMP-activated protein kinase in muscle during exercise. *Am J Physiol* 1996;270:E299–304.
  63. Motoshima H, Goldstein BJ, Igata M, Araki E. AMPK and cell proliferation-AMPK as a therapeutic target for atherosclerosis and cancer. *J Physiol* 2006;574:63–71.
  64. Xiang X, Saha AK, Wen R, Ruderman NB, Luo Z. AMP-activated protein kinase activators can inhibit the growth of prostate cancer cells by multiple mechanisms. *Biochem Biophys Res Commun* 2004;321:161–7.
  65. Imamura K, Ogura T, Kishimoto A, Kaminishi M, Esumi H. Cell cycle regulation via p53 phosphorylation by a 5'-AMP activated protein kinase activator, 5-aminoimidazole-4-carboxamide-1- $\beta$ -D-ribofuranoside, in a human hepatocellular carcinoma cell line. *Biochem Biophys Res Commun* 2001;287:562–7.
  66. Igata M, Motoshima H, Tsuruzoe K, et al. Adenosine monophosphate-activated protein kinase suppresses vascular smooth muscle cell proliferation through the inhibition of cell cycle progression. *Circ Res* 2005;97:837–44.
  67. Ruderman NB, Saha AK, Kraegen EW. Minireview: malonyl CoA, AMP-activated protein kinase, and adiposity. *Endocrinology* 2003;144:5166–71.
  68. Funabashi H, Kawaguchi A, Tomoda H, Omura S, Okuda S, Iwasaki S. Binding site of cerulenin in fatty acid synthetase. *J Biochem (Tokyo)* 1989;105:751–5.
  69. Weiss L, Hoffmann GE, Schreiber R, et al. Fatty-acid biosynthesis in man, a pathway of minor importance. Purification, optimal assay conditions, and organ distribution of fatty-acid synthase. *Biol Chem Hoppe Seyler* 1986;367:905–12.



the conclusion of the study, the control animals weighed  $97.5 \pm 1.2\%$  of pretreatment weight compared with  $95.0\% \pm 2.5\%$  for the C93-treated animals (not significant;  $P > 0.050$ , two-tailed  $t$  test). Thus, nonoptimized treatment of the SKOV3 xenograft with C93 produced a significant antitumor response without significant weight loss. Histopathologic analysis of C93 SKOV3 xenografts showed extensive tumor cytotoxicity characterized by acute inflammation (Fig. 4C, arrows) compared with vehicle control (Fig. 4D). Necropsy of the animals at the conclusion of both studies showed no gross or microscopic evidence of cytotoxicity to proliferating cell compartments including skin, gastrointestinal tract, or bone marrow (data not shown).


## Discussion

With the extensive array of human cancers that express high levels of FAS and the numerous studies documenting the cytotoxic effects of FAS inhibition both *in vitro* and *in vivo* (6), FAS holds promise as a new drug target for cancer therapy. Whereas our initial studies with C75, a synthetic FAS inhibitor (30), showed significant antitumor activity against human xenografts, treatment was limited by dramatic weight loss (18, 31, 56). We and our collaborators sought to understand the mechanism responsible for C75-induced weight loss to determine if it was (a) common to all FAS inhibitors, (b) unique to C75, (c) due to action at a second drug target, or (d) the result of nonspecific sickness behavior. It now seems that C75-induced weight loss occurred predominantly from increased fatty acid oxidation via stimulation of CPT-1, rather than solely through FAS inhibition (57). The pharmacologic separation

of CPT-1 stimulation from FAS inhibition led to the development of C93, which affords substantial anticancer cytotoxicity *in vitro* and *in vivo* without the weight loss seen with C75 treatment.

C93 also allowed the further pursuit of the cytotoxic mechanism of action of FAS inhibition. The involvement of AMPK in the hypothalamic response to C75 (37) and studies linking AMPK activation to cell cycle arrest in transformed cells (55) led us to hypothesize that AMPK may be involved in the cytotoxic response of FAS inhibition. AMPK is a heterotrimeric complex consisting of catalytic subunit  $\alpha 1$  or  $\alpha 2$  and regulatory  $\beta$  and  $\gamma$  subunits, which function as a serine/threonine kinase (58). Once activated by phosphorylation, AMPK inactivates anabolic biosynthetic enzymes for cholesterol synthesis, such as hydroxymethylglutaryl-CoA reductase, and for fatty synthesis, such as ACC, curbing ATP utilization (38, 52). Concomitantly, AMPK stimulates catabolic processes that enhance ATP production such as glucose uptake, glycolysis, and fatty acid oxidation (59–62).

AMPK itself has been proposed as a therapeutic target for cancer (55, 63, 64). AMPK activation has been shown to induce cell cycle arrest in transformed cell lines such as HepG2 (55, 65) and in nontransformed vascular smooth muscle cells (66). In a recent study, AICAR treatment reduced cell proliferation in a number of cancer cell lines *in vitro* including C<sub>6</sub> rat transformed glioma, MCF-7 human breast, and PC3 human prostate (55). AICAR produced an S-phase arrest likely mediated by p21, p27, and p53 proteins and inhibition of Akt activation. *In vivo* treatment of C<sub>6</sub> glioma cells in rats with AICAR reduced tumor mass and cell proliferation. Importantly, AICAR inhibited proliferation of *LKB*<sup>−/−</sup> mouse embryo fibroblasts showing independence of the *LKB*



**Figure 2.** C93 increased glucose oxidation and altered redox balance in SKOV3 cells. **A**, glucose oxidation was increased within 60 min of C93 treatment ( $P = 0.0009$ ). **B**, despite increased glucose oxidation, the NAD<sup>+</sup>/NADH redox couple declined ( $P < 0.0001$ ). **C**, NADPH levels increased 5 min after C93 treatment but began to decline after 2 h ( $P = 0.008$ ). **D**, the NADPH/NADP<sup>+</sup> redox couple was reduced 2 h after C93 treatment ( $P = 0.013$ ).  $P$  values were calculated using one-way ANOVA analysis;  $P$  values for all individual time points were calculated with time 0 as the control value using Dunnett's posttest (\*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ ; Prism 4.0, GraphPad Software).

cancer suppressor gene (55). Whereas these studies indicate a link between cancer cell growth and the status of AMPK activation, nonselective activation of AMPK in the whole organism may not be suitable for therapy due to the pleiotropic effects of AMPK activation (67).

FAS inhibition was first linked to AMPK activation through studies with C75 in neurons and the central nervous system (36, 37). In primary cultures of cortical neurons, C75 and cerulenin, a natural product FAS inhibitor (68), initially activated AMPK within minutes followed by inactivation over hours with no evidence of cellular injury. In addition, ACC inhibition with TOFA failed to show a change in AMPK activation (36). In the hypothalamus, C75 rapidly inactivated AMPK, leading to anorexia. AICAR reversed the effect of AMPK whereas compound C, an AMPK inhibitor, mimicked the effect of C75 (37). Thus, feeding behavior, FAS activity, and AMPK activation were involved in the hypothalamic control of food intake. Because C75 was shown to inhibit the growth of human cancer cells, we hypothesized that AMPK may also be involved in the cytotoxic effect of FAS inhibition on cancer cells (13, 18–21).

The increased NADPH levels following C93 treatment suggests that significant inhibition of FAS occurs within minutes of


treatment. We detected significant inhibition of the fatty acid synthesis pathway within 30 min of C93 treatment with the lag time likely due to the limitations of [ $^{14}$ C]acetate metabolic labeling of lipids. FAS inhibition was followed by a significant reduction in the AMP/ATP ratio also within 30 min of treatment. If an increase in the AMP/ATP ratio is biologically significant enough to signal an energy poor state, it should also activate AMPK through phosphorylation. Both by immunoblot and SAMS peptide assay, AMPK was activated within 30 min following C93 treatment. Thus, C93 rapidly inhibited FAS and induced an energy-poor state in the cancer cells with activated AMPK.

If the downstream signal transduction pathways are intact in SKOV3 cells, AMPK activation should reduce anabolic pathways and enhance catabolism. Two hours after C93 treatment, ACC was phosphorylated and fatty acid synthesis was further reduced, likely representing inhibition of ACC activity due to its phosphorylation. Within 60 min posttreatment, glucose oxidation was significantly increased. Thus, AMPK activation phosphorylated and inhibited ACC to reduce fatty acid synthesis while increasing catabolic glucose oxidation. Interestingly, AMPK activation has been shown to suppress FAS expression in primary cultured hepatocytes (53) and in androgen-independent (DU145 and PC3) and androgen-sensitive (LNCaP) cells (64). Whereas chronic AMPK activation may change the phenotype of cancer cells including FAS expression, it is unlikely that the rapidity of the changes following C93 treatment could be explained by reduced FAS expression because the half-life of the enzyme in cancer cells is ~12 h (data not shown). Indeed, over the course of these experiments, no changes in the FAS expression were identified by immunoblot.


Although the increased AMP/ATP ratio following C93 treatment activated AMPK and increased glucose oxidation, the AMP/ATP ratio continued to increase. Moreover, the oxidative potential of the cells reflected in the NAD<sup>+</sup>/NADH ratio declined steadily 30 min posttreatment. Thus, although glucose oxidation increased, it was unable to compensate for the loss of ATP. The declining NAD<sup>+</sup>/NADH ratio indicated a diminished capacity to further increase catabolism in response to the increasing AMP/ATP ratio. The anabolic redox couple NADP<sup>+</sup>/NADPH was also substantially affected by C93 treatment. Within 10 min of treatment, NADPH levels were significantly elevated, perhaps reflecting the rapid inhibition of FAS leading to substrate accumulation. By 60 min, however, the NADPH/NADP<sup>+</sup> ratio began decreasing, which could impede macromolecular synthesis. Indeed, potent inhibition of DNA replication and S-phase progression in human cancer cells was noted with both cerulenin and C75 treatment (24, 25).

If AMPK activation is responsible for triggering cytotoxicity, prior treatment with compound C to block AMPK activation should blunt the cytotoxic effect of FAS inhibition. Indeed, pretreatment with compound C substantially rescued SKOV3 cells from C93 treatment. AMPK inactivation with compound C had no significant cytotoxic effect on SKOV3 cells. These data implicate AMPK activation in the cytotoxic mechanism of action of pharmacologic FAS inhibition with C93.

Prior studies from our laboratory with human breast cancer cells showed that FAS inhibition was substantially more cytotoxic than ACC inhibition (18). These data suggested that substrate accumulation, not end-product (fatty acid) depletion, was responsible for the cytotoxicity. In SKOV3 cells, although the IC<sub>50</sub> for pathway inhibition with TOFA was >25-fold less than C93, the TOFA LC<sub>50</sub> for SKOV3 cells was 16-fold greater. Moreover, TOFA treatment did not lead to AMPK phosphorylation. C93 and AICAR, both AMPK



**Figure 3.** Compound C, an AMPK inhibitor, rescued SKOV3 cells from C93 cytotoxicity; TOFA, an ACC inhibitor, did not activate AMPK. **A**, SKOV3 cells pretreated with compound C were not susceptible to C93-induced cytotoxicity. **B**, immunoblot showed that treatment of SKOV3 cells with TOFA did not activate AMPK. **C**, both C93 and AICAR, AMPK activators, were significantly cytotoxic whereas TOFA failed to induce significant cytotoxicity compared with vehicle-treated controls (two-tailed *t* tests: \*, *P* < 0.05; \*\*, *P* < 0.01; \*\*\*, *P* < 0.001; Prism 4.0, GraphPad Software).



**Figure 4.** C93 inhibited the growth of SKOV3 xenografts. **A**, i.p. C93 treatment of an i.p. SKOV3 xenograft ( $n = 8$ ) showed significant reduction in tumor volume with three animals tumor-free (two-tailed  $t$  test; GraphPad Software). Three C93-treated animals died from traumatic i.p. injections. **B**, to assess systemic C93 efficacy, s.c. SKOV3 xenografts were treated with i.p. C93. There was substantial reduction in tumor growth (repeated measures two-way ANOVA; Prism 4.0 GraphPad Software). **C**, xenograft treated with C93. Note the geographic areas of intense intratumoral acute inflammation and tumor cytotoxicity (arrows;  $\times 200$ ), which are absent from the vehicle control (**D**;  $\times 200$ ; H&E staining).

activators, were substantially cytotoxic to SKOV3 cells after 24 h, with no cytotoxicity detected with TOFA.

To determine the efficacy and any potential toxicity of C93 *in vivo*, we treated i.p. and s.c. SKOV3 xenograft-bearing athymic mice with C93 in two pilot studies. In the i.p. SKOV3 xenograft model (20), C93 daily i.p. treatment (50 mg/kg) substantially reduced tumor growth compared with vehicle controls with no identifiable disease in three animals. We next studied s.c. SKOV3 xenografts, escalating the dose to twice-daily C93 treatment for 2 weeks. The C93-treated mice had a 3-fold reduction in tumor growth compared with vehicle controls. Histopathology of the xenograft 4 h after treatment showed evidence of ongoing cytotoxicity characterized by foci of acute inflammation. No cytotoxic effect on other proliferating cell compartments such as skin, gastrointestinal tract, or bone marrow was noted.

The pharmacologic inhibition of FAS with C93 causes rapid energy depletion, AMPK activation, and substantial cytotoxicity *in vitro*. Pilot studies in the SKOV3 xenograft showed significant antitumor effect without evidence of significant toxicity or weight loss. Because FAS is up-regulated in cancer cells but not in normal human tissues (69), pharmacologic FAS inhibition may provide a means to selectively activate AMPK and induce a cytotoxic response in ovarian cancer while sparing most of normal human tissues from the pleiotropic effects of AMPK activation.

## Acknowledgments

Received 9/25/2006; revised 12/13/2006; accepted 1/17/2007.

Grant support: NIH grants CA87850 (F.P. Kuhajda), Small Business Innovation Research National Cancer Institute grant 1R44CA99435 (to FASgen), and FASgen.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

## References

- Armstrong DK. Relapsed ovarian cancer: challenges and management strategies for a chronic disease. *Oncologist* 2002;7 Suppl 5:20-8.
- Bhoola S, Hoskins WJ. Diagnosis and management of epithelial ovarian cancer. *Obstet Gynecol* 2006;107:1399-410.
- Wakil S. Fatty acid synthase, a proficient multifunctional enzyme. *Biochemistry* 1989;28:4523-30.
- Alo PL, Visca P, Framarino ML, et al. Immunohistochemical study of fatty acid synthase in ovarian neoplasms. *Oncol Rep* 2000;7:1383-8.
- Gansler TS, Hardman W III, Hunt DA, Schaffel S, Hennigar RA. Increased expression of fatty acid synthase (OA-519) in ovarian neoplasms predicts shorter survival. *Hum Pathol* 1997;28:686-92.

- Kuhajda FP. Fatty acid synthase and cancer: new application of an old pathway. *Cancer Res* 2006;66:5977-80.
- Yang YA, Han WE, Morin PJ, Chrest FJ, Pizer ES. Activation of fatty acid synthesis during neoplastic transformation: role of mitogen-activated protein kinase and phosphatidylinositol 3-kinase. *Exp Cell Res* 2002;279:80-90.
- Yang YA, Morin PJ, Han WE, et al. Regulation of fatty acid synthase expression in breast cancer by sterol regulatory element binding protein-1c. *Exp Cell Res* 2003;282:132-7.
- Van de Sande T, De Schrijver E, Heyns W, Verhoeven G, Swinnen JV. Role of the phosphatidylinositol 3'-kinase/PTEN/Akt kinase pathway in the overexpression of fatty acid synthase in LNCaP prostate cancer cells. *Cancer Res* 2002;62:642-6.

- Bull JH, Ellison G, Patel A, et al. Identification of potential diagnostic markers of prostate cancer and prostatic intraepithelial neoplasia using cDNA microarray. *Br J Cancer* 2001;84:1512-9.
- Swinnen JV, Roskams T, Joniau S, et al. Overexpression of fatty acid synthase is an early and common event in the development of prostate cancer. *Int J Cancer* 2002;98:19-22.
- Epstein JI, Carmichael M, Partin AW. OA-519 (fatty acid synthase) as an independent predictor of pathologic state in adenocarcinoma of the prostate. *Urology* 1995;45:81-6.
- Pizer ES, Pflug BR, Bova GS, Han WE, Udan MS, Nelson JB. Increased fatty acid synthase as a therapeutic target in androgen-independent prostate cancer progression. *Prostate* 2001;47:102-10.
- Alo PL, Visca P, Marci A, Mangoni A, Botti C, Di